

Innovation. Quality. Solutions.

Kusabi® 300 SC FUNGICIDE

Reliable chemistry for powdery mildew programs

Kusabi contains 300 g/L pyriofenone (MOA Group 50)

The Product

Kusabi® 300 SC is a persistent fungicide for the control of powdery mildew in cucurbits.

Kusabi contains the active ingredient pyriofenone. It is the second product to be commercialised from the actin disruption (Group 50) chemical group.

Actin provides strength to the outer structure of fungi, and in its absence, powdery mildew cannot enter the plant or grow normally.

Kusabi has strong vapour activity which supports optimal spray coverage.

Application Rate

Kusabi is registered at 300 or 500 mL/ha. Use the higher rate when disease pressure is high. Apply in a spray program with other fungicides at 7 to 10 day spray intervals.

Kusabi should be applied at 500 mL/ha in a medium to high water volume (500 to 1000 L/ha) as a protectant treatment for crops grown in areas where powdery mildew occurs frequently and disease levels are normally high.

Kusabi can be used at 300 mL/ha when powdery mildew pressure is normally at lower levels, or when mixed with Du-Wett adjuvant (applied at 200 mL/ha). The use of Du-Wett will also assist spray coverage, especially when lower water volumes (<500 L/ha) are used.

Safety


Kusabi has an excellent toxicological and environmental profile when used as directed. In a trial to assess the safety of the beneficial species green lacewing, Persimilis and Trichogramma, Kusabi was not acutely toxic. Kusabi has also been assessed by the APVMA as not toxic to bees.

Timing

Kusabi is a new fungicide for the control of powdery mildew in cucurbits. Kusabi performs best when applied as a protectant fungicide to a clean crop. An excellent partner to rotate with other fungicides, Kusabi is ideally suited to follow Flute® in a fungicide spray program from flowering.

Field Data

Field trials across Australia have shown Kusabi to be highly effective for the control of powdery mildew.



Summary

- 1. Kusabi** is a powdery mildew fungicide developed and manufactured by ISK, an R&D company from Japan.
- 2. Kusabi** is a single-site fungicide that is subject to a resistance management strategy. This limits the number of applications to three per crop, with strict alternation of chemical groups.
- 3. Rainfast:** 1 hour.
- 4. Compatibility:** Kusabi has demonstrated physical compatibility with a wide range of commonly used insecticides and fungicides,

however not all products have yet been tested. For updated information, please refer to agnova.com.au

- 5. Crop safety:** Excellent crop safety when used as directed.
- 6.** Apply to a clean crop after the application of a fungicide such as Flute, as part of a complete disease control program.
- 7. WHP:** Not required when used as directed.

Innovation. Quality. Solutions.

Kusabi® 300 SC FUNGICIDE

Resistance Management Strategy for Cucurbits:

Group 3: (DMI, pyrimidine)

Group 7: SDHI (Succinate dehydrogenase inhibitors)

Group 8: (Hydroxy-(2-amino-) pyrimidine)

Group U6: (Phenyl-acetamide)

Group 11: (Quinone outside Inhibitor) and other "systemic" fungicides

Group 13: Aza-naphthalenes

Group 50: (Actin Inhibitor) fungicides

- 1. Start disease control early.** DO NOT wait for powdery mildew to appear before spraying, but start as soon as practicable after crop emergence.
- 2. Use protectant sprays in early crop growth.** Apply protectant sprays up to the fruit set stage of the crop if the disease normally occurs during this period. If this schedule is interrupted (e.g. by rain) use a tank mix of protectant plus systemic before recommencing the protectant program.
- 3. After fruit set, use systemic fungicides in one or more of the following ways:**
 - a. Tank mix systemic fungicides with a protectant fungicide AND use fungicides from at least two different systemic activity groups per crop.
 - b. Alternate systemic fungicides with a protectant fungicide AND use fungicides from at least two different systemic activity groups per crop.
 - c. Alternate systemic fungicides from at least three different activity groups per crop.
 - d. Apply Group 11 fungicides preventatively.
- 4. Use a maximum of one Group 11 containing spray out of every three fungicide applications.**
- 5. DO NOT use consecutive applications of Group 11 or Group U6 fungicides.**
- 6. DO NOT apply more than two Group 11 or Group U6 products per crop.**
- 7. DO NOT apply more than three Group 7 or Group 13 products per crop and no more than two consecutive applications per year**



CROP STAGE						
DOMINANT DISEASE	ESTABLISHMENT	FLOWERING		FRUIT SET		FRUIT MATURITY
Powdery mildew	Group 3, 7, (7+11), 11 or 13	Group 3, 7, (7+11), 11 or 13	FLUTE (U6)	KUSABI (50)	FLUTE (U6)	KUSABI (50)

The information provided herein may include extracts from the product label and does not constitute the complete directions for use. READ THE PRODUCT LABEL THOROUGHLY BEFORE OPENING OR USING KUSABI

AgNova Technologies Pty Ltd
ABN 70 097 705 158
PO Box 2069 Box Hill North
Victoria 3129 Australia
Ph 03 9899 8100

Data referred to herein were generated using products and rates that were registered at the time. Always read and follow product labels.

AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.

© Copyright AgNova Technologies 2017. KUSCUC190913

© Kusabi is a registered trademark of Ishihara Sangyo Kaisha Ltd. Japan. Flute is a registered trademark of Nippon Soda Co. Ltd.

© Du-Wett is a registered trademark of Elliott Chemicals Limited used under licence by Nufarm Australia Limited

* Registered trademark